

Contents

Mitochondrial genetics

- 429** Brain imaging in mitochondrial respiratory chain deficiency: combination of brain MRI features as a useful tool for genotype/phenotype correlations *M Bricout, D Grévent, A S Lebre, M Rio, I Desguerre, P De Lonlay, V Valayannopoulos, F Brunelle, A Rötig, A Munnich, N Boddaert*

Disease loci

- 436** *ADAP2* in heart development: a candidate gene for the occurrence of cardiovascular malformations in NF1 microdeletion syndrome *M Venturin, S Carra, G Gaudenzi, S Brunelli, G R Gallo, S Moncini, F Cotelli, P Riva*
- 444** A missense mutation in the splicing factor gene *DHX38* is associated with early-onset retinitis pigmentosa with macular coloboma *M Ajmal, M I Khan, K Neveling, Y M Khan, M Azam, N K Waheed, C P Hamel, T Ben-Yosef, E De Baere, R K Koenekoop, R W J Collin, R Qamar, F P M Cremers*

Complex traits

- 449** Haplotypes of the inducible nitric oxide synthase gene are strongly associated with exhaled nitric oxide levels in adults: a population-based study *S Dahgam, L Modig, Å T Naluai, A-C Olin, F Nyberg*

Somatic mosaicism

- 455** Early postzygotic mutations contribute to de novo variation in a healthy monozygotic twin pair *G M Dal, B Ergüner, M S Sağıroğlu, B Yükesel, O E Onat, C Alkan, T Özçelik*

New loci

- 460** A homozygous nonsense CEP250 mutation combined with a heterozygous nonsense C2orf71 mutation is associated with atypical Usher syndrome *S Khateb, L Zelinger, L Mizrahi-Meissonnier, C Ayuso, R K Koenekoop, U Laxer, M Gross, E Banin, D Sharon*

- 470** Mutations in the enzyme glutathione peroxidase 4 cause Sedaghatian-type spondylometaphyseal dysplasia *A C Smith, A J Mears, R Bunker, A Ahmed, M MacKenzie, J A Schwartzenruber, C L Beaulieu, E Ferretti, FORGE Canada Consortium, J Majewski, D E Bulman, F C Celik, K M Boycott, G E Graham*
- 475** A novel stop mutation in the vascular endothelial growth factor-C gene (*VEGFC*) results in Milroy-like disease *E Balboa-Beltran, M J Fernández-Seara, A Pérez-Muñizuri, R Lago, C García-Magán, M L Couce, B Sobrino, J Amigo, A Carracedo, F Barros*

Phenotypes

- 479** Prediction of the age at onset in spinocerebellar ataxia type 1, 2, 3 and 6 *S T du Moncel, A Durr, M Rakowicz, L Nanetti, P Charles, A Sulek, C Mariotti, R Rola, L Schols, P Bauer, I Dufaure-Garé, H Jacobi, S Forlani, T Schmitz-Hübsch, A Filla, D Timmann, B P van de Warrenburg, C Marelli, J-S Kang, P Giunti, A Cooke, L Baliko, M Bela, S Boesch, S Szymanski, J Berciano, J Infante, K Buerk, M Masciullo, R D Fabio, C Depondt, S Ratka, G Stevanin, T Klockgether, A Brice, J-L Golmard*

Cognitive and behavioural genetics

- 487** Involvement of the kinesin family members *KIF4A* and *KIF5C* in intellectual disability and synaptic function *M H Willemsen, W Ba, W M Wissink-Lindhout, A P M de Brouwer, S A Haas, M Bieneke, H Hu, L E L M Vissers, H van Bokhoven, V Kalscheuer, N N Kasri, T Kleefstra*

PostScript

- 478** Corrections



Cover credit: Detecting post-zygotic SNVs in monozygotic twins, from Dal *et al.* pg 455.

Receive regular table of contents by email. Register using this QR code.



This article has been chosen by the Editor to be of special interest or importance and is freely available online.



This article has been made freely available online under the BMJ Journals Open Access scheme. See <http://jmg.bmj.com/site/about/guidelines.xhtml#open>



This journal is a member of and subscribes to the principles of the Committee on Publication Ethics <http://publicationethics.org/>



[MCQs] The online version of this article contains multiple choice questions hosted on BMJ Learning.